



IJM – Italian Journal of Mycology, Vol. 54 (2025): 1-11

<https://doi.org/10.6092/issn.2531-7342/19781>



Supplementary material

Squalene in fruiting bodies of true mushrooms changes in the process of vegetation


Ivan M. Korabel¹, Lidiia V. Panchak¹, Alina R. Zyn², Volodymyr O. Antonyuk^{1,3}

¹Department of Pharmaceutical, Organic and Bioorganic Chemistry, Faculty of Pharmacy, Danylo Halytsky Lviv National Medical University, Pekarska St., 69, 79010, Ukraine.

²Lviv Research Forensic Centre of MIA of Ukraine, Koniushynna St., 24, 79000, Ukraine.

³Department of Regulation of Cell Proliferation and Apoptosis, Institute of Cell Biology NAS of Ukraine, Drahomanov St. 14/16, 79005, Lviv, Ukraine.

Corresponding author e-mail: antonyukvo@gmail.com



Supplementary Fig. S1 - Thin-layer chromatogram of lipophilic fractions of fungi and squalene.

Numbers indicate the tracks of lipophilic extracts: 1 - *Mucidula múcida*, 2 - *Lycoperdon pyriforme*, 3 - *Laetiporus sulphureus*, 4 - *Tyromyces chioneus*, 5 - *Sarcoscypha coccinea*, 6 - fractions 1-4 of the lipophilic extract of *Sarcoscypha coccinea* (Fig. 1), 7 - *Amaranthus caudatus*, 8 - squalene from Sigma-Aldrich.