

IJM – Italian Journal of Mycology, Vol. 53 (2024): 160-165

https://doi.org/10.6092/issn.2531-7342/19902



## Supplementary material

## Medicinal wood decay fungi on large old oaks in the Madonie Mountains (Sicily)

Fortunato Cirlincione<sup>1</sup>, Giuseppe Venturella<sup>2</sup>, Maria Letizia Gargano<sup>1</sup>, Giulia Mirabile<sup>3</sup>, Gaetano Balenzano<sup>1</sup>, Raimondo Pardi<sup>1</sup>, Valeria Ferraro<sup>4</sup>


Corresponding author e-mail: <a href="mailto:giumirabile@gmail.com">giumirabile@gmail.com</a>


<sup>&</sup>lt;sup>1</sup>Department of Soil, Plant, and Food Sciences, University of Bari "Aldo Moro"; Via G. Amendola n. 165/A, 70126 Bari, Italy

<sup>&</sup>lt;sup>2</sup>Department of Agricultural, Food and Forest Sciences, University of Palermo; Viale delle Scienze Bldg. 5, 90128 Palermo, Italy

<sup>&</sup>lt;sup>3</sup>Provincial Agriculture Inspectorate of Palermo; Via C. Camalliani n. 54, 90145 Palermo, Italy

<sup>&</sup>lt;sup>4</sup>Department of Pharmacy – Pharmaceutical Sciences, University of Bari "Aldo Moro", University Campus "Ernesto Quagliarello"; Via E. Orabona n.4, 70125 Bari, Italy





**Supplementary Fig. S1** – ITS-Phylogenetic tree based on Neighbour-joining analysis a) *Laetiporus* spp. and b) *Pseudoinonotus* and *Inonotus* spp. Bootstrap values are indicated at nodes. Black triangles indicate the sequences obtained in this survey and deposited in GenBank database. *Fomitopsis pinicola* (Sw.) P. Karst. and *Fomitiporia mediterranea* M. Fisch. were used as outgroups. Bootstrap test were performed with 1000 replications.